arXiv:math/0407189v1 [math.AT] 11 Jul 2004

CELLULAR COCHAIN ALGEBRAS AND TORUS ACTIONS

ILIA V. BASKAKOV, VICTOR M. BUCHSTABER, AND TARAS E. PANOV

We prove that the *integral* cohomology algebra of the moment-angle complex $\mathcal{Z}_K$ [1] is isomorphic to the Tor-algebra of the face ring of simplicial complex K. The proof relies upon the construction of a cellular approximation of the diagonal map $\Delta\colon \mathcal{Z}_K \to \mathcal{Z}_K \times \mathcal{Z}_K$. Cellular cochains do not admit a functorial associative multiplication because a proper cellular diagonal approximation does not exist in general. The construction of moment-angle complexes is a functor from the category of simplicial complexes to the category of spaces with torus action. We show that in this special case the proposed cellular approximation of the diagonal is associative and functorial with respect to those maps of moment-angle complexes which are induced by simplicial maps.

The *face ring* of a complex K on the vertex set $[m] = \{1, \ldots, m\}$ is the graded quotient ring $\mathbb{Z}[K] = \mathbb{Z}[v_1, \ldots, v_m]/(v_\omega\colon \omega \notin K)$ with $\deg v_i = 2$ and $v_\omega = v_{i_1} \cdots v_{i_k}$, where $\omega = \{i_1, \ldots, i_k\} \subseteq [m]$. Let BT^m be the classifying space for the m-dimensional torus, endowed with the standard cell decomposition. Consider a cellular subcomplex $DJ(K) := \bigcup_{\sigma\in K} BT^\sigma \subseteq BT^m$, where $BT^\sigma = \{x = (x_1, \ldots, x_m) \in BT^m : x_i = pt \text{ for } i \notin \sigma\}$. Using the cellular decomposition we establish a ring isomorphism $H^*(DJ(K)) \cong \mathbb{Z}[K]$ (see [2, Lemma 2.8]). Let $D^2 \subset \mathbb{C}$ be the unit disc and set $B_\omega := \{(z_1, \ldots, z_m) \in (D^2)^m : |z_j| = 1 \text{ for } j \notin \omega\}$. The *moment-angle complex* is the T^m-invariant subspace $\mathcal{Z}_K := \bigcup_{\sigma\in K} B_\sigma \subseteq (D^2)^m$. As it is shown in [1, Ch. 6], the spaces $DJ(K)$ and $\mathcal{Z}_K$ are homotopy equivalent to the spaces introduced in [3], which justifies our notation. Complexes $\mathcal{Z}_K$ provide an important class of torus actions. The space $\mathcal{Z}_K$ is the homotopy fibre of the inclusion $DJ(K) \hookrightarrow BT^m$; it appears also as the level surface of the moment map used in the construction of *toric varieties* via symplectic reduction; it is also homotopy equivalent to the complement of the *coordinate subspace arrangement* determined by K, see [1, § 8.2].

Theorem 1. *There is a functorial in K isomorphism of algebras*

$$H^*(\mathcal{Z}_K;\mathbb{Z}) \cong H\bigl[\Lambda[u_1, \ldots, u_m] \otimes \mathbb{Z}[K], d\bigr] \cong \operatorname{Tor}_{\mathbb{Z}[v_1,\ldots,v_m]}\bigl(\mathbb{Z}[K], \mathbb{Z}\bigr),$$

where the algebra in the middle is the cohomology of the differential graded algebra with $\deg u_i = 1$, $\deg v_i = 2$, $du_i = v_i$, $dv_i = 0$.

In the case of rational coefficients this theorem was proved in [4] using spectral sequences techniques (see also [1, Th. 7.6, Probl 8.14]). Our new proof uses a construction of cellular cochain algebra. Another proof of Theorem 1 follows from a recent independent work of M. Franz [5, Th. 1.2].

The work is supported by the Russian Foundation for Basic Research, grant no. 02-01-00659, and by the Leading Scientific School support, grant no. 2185.2003.1.

Proof of Theorem 1. We prove just the first isomorphism, since the second is a standard consequence of the Koszul resolution (see [1] for details). Introduce an extra grading by setting $\operatorname{bideg} u_i = (-1, 2)$, $\deg v_i = (0, 2)$, and consider a quotient algebra

$$R^*(K) := \Lambda[u_1, \dots, u_m] \otimes \mathbb{Z}[K]/(v_i^2 = u_i v_i = 0,\ i = 1, \dots, m).$$

Let $\varrho\colon \Lambda[u_1, \dots, u_m] \otimes \mathbb{Z}[K] \to R^*(K)$ be the canonical projection. Using the finite additive basis in $R^*(K)$ consisting of monomials of the type $u_\omega v_\sigma$, where $\omega \subseteq [m]$, $\sigma \in K$ and $\omega \cap \sigma = \varnothing$, we define an additive inclusion $\iota\colon R^*(K) \to \Lambda[u_1, \dots, u_m] \otimes \mathbb{Z}[K]$ which satisfies $\varrho \cdot \iota = \mathrm{id}$. We claim that ϱ induces an isomorphism in the cohomology. To see this we construct a cochain homotopy operator s such that $ds + sd = \mathrm{id} - \iota \cdot \varrho$. If $K = \varDelta^{m-1}$ (the full simplex) the algebra $\Lambda[u_1, \dots, u_m] \otimes \mathbb{Z}[K]$ is $E = E_m = \Lambda[u_1, \dots, u_m] \otimes \mathbb{Z}[v_1, \dots, v_m]$, and $R^*(\varDelta^{m-1})$ is isomorphic to the algebra $R^*(\varDelta^0)^{\otimes m}$, where $R^*(\varDelta^0) = \Lambda[u] \otimes \mathbb{Z}[v]/(v^2 = uv = 0)$. A direct calculation shows that for $m = 1$ the map $s_1\colon E^{0,*} = \mathbb{Z}[v] \to E^{-1,*}$ given by $s_1(a_0 + a_1 v + \dots + a_j v^j) = (a_2 v + a_3 v^2 + \dots + a_j v^{j-1})u$ is the required cochain homotopy. Then we may assume by induction that for $m = k - 1$ there is a cochain homotopy operator $s_{k-1}\colon E_{k-1} \to E_{k-1}$. Since $E_k = E_{k-1} \otimes E_1$, $\varrho_k = \varrho_{k-1} \otimes \varrho_1$ and $\iota_k = \iota_{k-1} \otimes \iota_1$, the map $s_k = s_{k-1} \otimes \mathrm{id} + \iota_{k-1}\varrho_{k-1} \otimes s_1$ is a cochain homotopy between id and $\iota_k \varrho_k$, which finishes the proof for $K = \varDelta^{m-1}$. In the case of arbitrary complex K the algebras $\Lambda[u_1, \dots, u_m] \otimes \mathbb{Z}[K]$ and $R^*(K)$ are obtained from E_m and $R^*(\varDelta^{m-1})$ respectively by factoring out a monomial ideal. This factorisation does not affect the properties of the operator s, which therefore establishes the required cochain homotopy.

Consider a cellular subdivision of the polydisc $(D^2)^m$ with each D^2 subdivided into the cells 1, T and D of dimensions 0, 1 and 2 respectively. Therefore, the cells of $(D^2)^m$ are encoded by words $\mathcal{T} \in \{D, T, 1\}^m$. Assign to each pair of subsets $\sigma, \omega \subseteq [m]$, $\sigma \cap \omega = \varnothing$ the word $\mathcal{T}(\sigma, \omega)$ which has the letter D on the positions numbered by σ and letter T on the positions with numbers from ω. By the construction, the word $\mathcal{T}(\sigma, \omega)$ corresponds to a cell from $\mathcal{Z}_K \subset (D^2)^m$ if and only if $\sigma \in K$. The cellular cochain complex $C^*(\mathcal{Z}_K)$ has an additive basis consisting of the cochains of the type $\mathcal{T}(\sigma, \omega)^*$. It follows that the dimensions of the graded components of $C^*(\mathcal{Z}_K)$ and $R^*(K)$ coincide. Moreover, the map $g\colon R^*(K) \to C^*(\mathcal{Z}_K)$, $u_\omega v_\sigma \mapsto \mathcal{T}(\sigma, \omega)^*$, is an isomorphism of differential graded modules and induces an additive isomorphism $H[R^*(K)] \cong H^*(\mathcal{Z}_K)$. To finish the proof we construct a cellular approximation $\widetilde{\Delta}_K$ of the diagonal map $\Delta\colon \mathcal{Z}_K \to \mathcal{Z}_K \times \mathcal{Z}_K$ such that the multiplication induced on the cellular cochains coincides with the multiplication in $R^*(K)$ via the isomorphism g. For $K = \varDelta^0$ we have $\mathcal{Z}_K = D^2$. Set $z = \rho e^{i\varphi} \in D^2$ and define a map $\widetilde{\Delta}\colon D^2 \to D^2 \times D^2$ by

$$\widetilde{\Delta}(z) = \begin{cases} (1 + \rho(e^{2i\varphi} - 1), 1) & \text{for } \varphi \in [0, \pi], \\ (1, 1 + \rho(e^{2i\varphi} - 1)) & \text{for } \varphi \in [\pi, 2\pi). \end{cases}$$

This is a cellular map sending ∂D^2 to $\partial D^2 \times \partial D^2$ and homotopic to Δ in the class of such maps. For the corresponding multiplication in cellular cochains there is a multiplicative isomorphism $R^*(\varDelta^0) \to C^*(D^2)$. Passing to the case $K = \varDelta^{m-1}$, we see that the map $\widetilde{\Delta}\colon (D^2)^m \to (D^2)^m \times (D^2)^m$

gives rise to a multiplicative isomorphism

$$f\colon R^*(\mathit{\Delta}^{m-1}) = \Lambda[u_1,\dots,u_m]\otimes\mathbb{Z}[v_1,\dots,v_m]\big/(v_i^2 = u_iv_i = 0) \longrightarrow C^*\bigl((D^2)^m\bigr).$$

As it follows from the construction of $\mathcal{Z}_K$, the restriction of the map $\widetilde{\Delta}$ to $\mathcal{Z}_K$ is a cellular map $\widetilde{\Delta}_K\colon \mathcal{Z}_K \to \mathcal{Z}_K \times \mathcal{Z}_K$. Therefore, there is a multiplicative map $q\colon C^*((D^2)^m) \to C^*(\mathcal{Z}_K)$. Consider the commutative diagram

$$\begin{tikzcd}
R^*(\mathit{\Delta}^{m-1}) \arrow[r, "f"] \arrow[d, "p"'] & C^*\bigl((D^2)^m\bigr) \arrow[d, "q"] \\
R^*(K) \arrow[r, "g"] & C^*(\mathcal{Z}_K)
\end{tikzcd}.$$

Using the fact that p, f and q are ring homomorphisms, and p is an epimorphism, we deduce that g is also a ring homomorphism. Thus, g is a multiplicative isomorphism. □

References

[1] Victor M Buchstaber and Taras E Panov. *Torus Actions and Their Applications in Topology and Combinatorics.* Volume 24 of *University Lecture Series*, Amer. Math. Soc., Providence, R.I., 2002.

[2] Victor M Buchstaber and Taras E Panov. Torus actions, equivariant moment-angle complexes, and configurations of coordinate subspaces. (Russian) *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)*, 266:29–50, 2000. Translation in *J. Math. Sci.*, 113(4):558–568, 2003.

[3] Michael W Davis and Tadeusz Januszkiewicz. Convex polytopes, Coxeter orbifolds and torus actions. *Duke Math. J.*, 62(2):417–451, 1991.

[4] Victor M Buchstaber and Taras E Panov. Torus actions and the combinatorics of polytopes. (Russian) *Tr. Mat. Inst. Steklova*, 225:96–131, 1999. Translation in *Proc. Steklov Inst. Math.*, 2(225):87–120, 1999.

[5] Matthias Franz. On the integral cohomology of smooth toric varieties. Preprint, arXiv:math.AT/0308253, 2003.

Faculty of Mathematics and Mechanics, Moscow State University

E-mail address: `tpanov@mech.math.msu.su`